

KINVET BroodWell

“Vigorous Chicks, Vibrant Growth, Valued Returns.”

How to Use KINVET BroodWell

Daily Dosage for Chicks:

- Broilers: 0.5–1 ml per litre of drinking water
- Layers: 1–2 ml per litre of drinking water
- Note: Shake well before use

When to Give an Extra Boost:

- During periods of stress (e.g., vaccination, heat, transport)
 - If you observe signs like poor growth, ruffled feathers, or lethargy
 - Following antibiotic treatments to aid recovery
 - In cases of high mortality or uneven flock development.
-

Farmhand Favorites: Do's

1. Store It Right

Keep the bottle sealed in a cool, dry place away from direct sunlight and moisture

2. Mix Thoroughly

Ensure KINVET BroodWell is evenly mixed in the drinking water to provide uniform intake

3. Monitor Your Chicks

Regularly check for improvements in activity levels, feather quality, and growth rates

4. Maintain Clean Water Lines

Clean drinkers daily to prevent microbial growth and ensure effective delivery

Farmer's Don'ts – Avoid These Mistakes:

1. Don't Overdose

Exceeding the recommended dosage won't accelerate results and may be wasteful

2. Don't Leave Bottle Open

Exposure to air and humidity can degrade the product's efficacy

3. Don't Neglect Water Quality

Contaminated or stale water can hinder the effectiveness of KINVET BroodWell

4. **Don't Skip Hygiene Practices**

Dirty coops and equipment can stress chicks and compromise their health

5. **Don't Ignore Biosecurity Measures**

Implement footbaths, restrict visitor access, and disinfect equipment to prevent disease outbreaks.

Pro Tip

“Incorporate KINVET BroodWell into your chicks' daily routine, and witness a transformation in their vitality and growth—naturally enhancing their development without the need for antibiotics.”