

# KINVET VaxiBlue

“See the vaccine. Protect every bird.”

## How to Use KINVET VaxiBlue

- **Usage:** Mix 10 g KINVET VaxiBlue per 100 litre of water, or as directed by your poultry nutritionist or veterinarian.

### Method of Administration:

- Dissolve thoroughly in water before mixing in the vaccine.
- Ensure uniform distribution across all water lines or drinkers.

### Frequency of Use:

- Use only during vaccination via the drinking water route.
  - Recommended for each live vaccine cycle (ND, IB, IBD, etc.).
- 

## Farmer & Vet Practices – Do's & Don'ts

### DO's

- Use clean containers and fresh water to dissolve KINVET VaxiBlue.
- Administer within 30 minutes of mixing with vaccine for best results.
- Observe birds during vaccination – blue tongue stains confirm intake.
- Flush water lines before and after vaccination to avoid residue build-up.
- Store in a cool, dry place away from sunlight and moisture.
- Keep pack tightly sealed after each use to maintain efficacy.
- Consult your vet before adding to vaccine protocols for large flocks.

### DON'Ts

- Don't use KINVET VaxiBlue as a general-purpose water conditioner.
- Don't mix with antibiotics, acids, or unrelated supplements.
- Don't exceed recommended dosage – more dye doesn't mean better results.

- ✘ Don't store diluted solution – prepare fresh before each use.
- ✘ Don't rely solely on guesswork—blue water equals assurance.

**Pro Tip**

*“Birds won't tell you they missed a dose – but their **tongues will.**”*

Use KINVET VaxiBlue to catch under-vaccination before it becomes a flock-wide issue.